


WORLD DATA CENTER A

Upper Atmosphere Geophysics



**SYNOPTIC RADIO MAPS OF THE SUN AT 3.3mm
1970-1973**

1



May 1978

WORLD DATA CENTER A
National Academy of Sciences
2101 Constitution Avenue, N.W.
Washington, D.C., U.S.A., 20418

World Data Center A consists of the Coordination Office

and seven Subcenters:

World Data Center A
Coordination Office
National Academy of Sciences
2101 Constitution Avenue, N.W.
Washington, D.C., U.S.A., 20418
[Telephone: (202) 389-6478]

Glaciology:

World Data Center A: Glaciology
[Snow and Ice]
Inst. of Arctic & Alpine Research
University of Colorado
Boulder, Colorado, U.S.A. 80309
[Telephone: (303) 492-5171]

Meteorology (and Nuclear Radiation):

World Data Center A: Meteorology
National Climatic Center
Federal Building
Asheville, North Carolina, U.S.A. 28801
[Telephone: (704) 258-2850]

Oceanography:

World Data Center A: Oceanography
National Oceanic and Atmospheric
Administration
Washington, D.C., U.S.A. 20235
[Telephone: (202) 634-7249]

Rockets and Satellites:

World Data Center A: Rockets and
Satellites
Goddard Space Flight Center
Code 601
Greenbelt, Maryland, U.S.A. 20771
[Telephone: (301) 982-6695]

NOTES:

1. World Data Centers conduct international exchange of geophysical observations in accordance with the principles set forth by the International Council of Scientific Unions. WDC-A is established in the United States under the auspices of the National Academy of Sciences.
2. Communications regarding data interchange matters in general and World Data Center A as a whole should be addressed to: World Data Center A, Coordination Office (see address above).
3. Inquiries and communications concerning data in specific disciplines should be addressed to the appropriate subcenter listed above.

Rotation of the Earth:

World Data Center A: Rotation
of the Earth
U.S. Naval Observatory
Washington, D.C., U.S.A. 20390
[Telephone: (202) 254-4023]


*Solar-Terrestrial Physics (Solar and
Interplanetary Phenomena, Ionospheric
Phenomena, Flare-Associated Events,
Geomagnetic Variations, Magnetospheric
and Interplanetary Magnetic Phenomena,
Aurora, Cosmic Rays, Airglow):*

World Data Center A
for Solar-Terrestrial Physics
Environmental Data Service, NOAA
Boulder, Colorado, U.S.A. 80303
[Telephone: (303) 499-1000, Ext. 6467]

*Solid-Earth Geophysics (Seismology,
Tsunamis, Gravimetry, Earth Tides,
Recent Movements of the Earth's
Crust, Magnetic Measurements,
Paleomagnetism and Archeomagnetism,
Volcanology, Geothermics):*

World Data Center A
for Solid-Earth Geophysics
Environmental Data Service, NOAA
Boulder, Colorado, U.S.A. 80303
[Telephone: (303) 499-1000, Ext. 6521]

WORLD DATA CENTER A for Solar-Terrestrial Physics



REPORT UAG-66

SYNOPTIC RADIO MAPS OF THE SUN AT 3.3mm 1970-1973

by

Earle B. Mayfield
Space Sciences Laboratory

and


Fred I. Shimabukuro
Electronics Research Laboratory

The Ivan A. Getting Laboratories
The Aerospace Corporation
El Segundo, California, USA 90245

May 1978


Published by World Data Center A for
Solar-Terrestrial Physics, NOAA, Boulder, Colorado
and printed by

U.S. DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
ENVIRONMENTAL DATA SERVICE
Asheville, North Carolina, USA 28801



SUBSCRIPTION PRICE: \$25.20 a year; \$17.30 additional for foreign mailing; single copy price varies.*
Checks and money orders should be made payable to the Department of Commerce, NOAA/NGSDC.
Remittance and correspondence regarding subscriptions should be sent to the National Geophysical and
Solar-Terrestrial Data Center, NOAA, Boulder, CO 80303.

*PRICE THIS ISSUE .75 CENTS



SYNOPTIC RADIO MAPS OF THE SUN AT 3.3 mm, 1970-1973

by

Earle B. Mayfield
Space Sciences Laboratory

and

Fred I. Shimabukuro
Electronics Research Laboratory

The Ivan A. Getting Laboratories
The Aerospace Corporation
El Segundo, California 90245

INTRODUCTION

The Aerospace Corporation millimeter radio telescope, which was completed in 1963, has been used extensively for investigations in planetary, solar, and stellar research. It is located in El Segundo, California, at $30^{\circ}54'53''N$ and $118^{\circ}22'30''W$ ($7h\ 53m\ 30s$) at a height of 38 m. Maps of the whole solar disk at 90 GHz (3.3-mm wavelength) were obtained routinely on a near daily basis from 1966 through 1973. In October 1973, loss of funding resulted in the discontinuation of the solar program. Since then, only infrequent observations of the Sun have been made to calibrate the radiometer.

The format used in presenting millimeter radio observations is the same as that employed for previous results [Mayfield et al., 1974] and is based on investigations of variations in the 3.3-mm emission and its correlation with photospheric magnetic fields, faculae, and centers of activity. Furthermore, the regions of enhanced millimeter emission are well-defined, stable features that can be readily identified and followed during their disk passage. The relatively infrequent, transient phenomena associated with flares, which have been discussed by Shimabukuro [1968], usually lie at the resolution limit of the antenna. As a result, synoptic maps are considered not only the most informative presentation but the most concise. The authors can provide separate maps for those individuals who require more detailed data or individual maps for specific days.

INSTRUMENT DESCRIPTION

The polar-mounted antenna is Cassegrain in design and 4.57 m (15 ft) in diameter [Jacobs and King, 1965; King et al., 1966]. Although the surface has been finished for operation at 400 GHz (1 mm), it is usually operated at 90 GHz (3.3 mm). At this wavelength the beamwidth at the half-power points is about 2.8 arc-min. Pointing accuracy and stability for the instrument is about 20 arc-sec. An on-line digital computer controls the antenna through a closed loop servo that permits a variety of tracking modes, including a mapping mode for solar studies. The radiometer is of the Dicke type in which switching is done at 465 Hz between the on-axis beam and a wide-beam sky temperature reference horn. The first stage of the radiometer is a single-end gallium arsenide crystal mixer that provides for frequency conversion to a 3-GHz intermediate amplifier. The system sensitivity for a signal-to-noise ratio of 1 at the output is about 0.6 K for a 1-second integration time.

For the results reported here, the computer-controlled antenna was programmed to obtain a square array of emission temperatures centered on the solar disk. A 19×19 matrix was obtained by rastering the antenna between adjacent points. A 4-second settle time was allowed between readings with a 1-second integration time for temperature measurement. These individual values were normalized to an undisturbed region near the center of the disk to eliminate the requirement of absolute data calibration. This reference temperature used for normalization was selected by analyzing millimeter observations previously obtained and by using the Fraunhofer Institute maps to select a quiet region free of plage or other disturbed emission. The finite beamwidth of the antenna at the solar limb restricts interpretation and thus the isotherms to regions less than $0.7R_{\odot}$ (normalized solar radius). The maps contain this limitation. Others include occasional pointing errors, typically greater than 20 arc-sec but less than 1 arc-min, that occur in centering the temperature matrix on the disk and slight differences in the emission temperatures owing to variations in cloud cover.

Figure 1 shows a typical map dated 30 June 1971 that has been superposed on an $H\alpha$ picture. Isotherms have been constructed on the temperature matrix at levels of 2% enhancement, and they indicate in this case the presence of two regions of significant enhancement--ones that can be identified with $H\alpha$ plage in centers of activity associated with spot group magnetic fields. The maximum enhancement in these millimeter regions usually occurs near the neutral line of a primary bipolar magnetic feature and shows daily variations associated with changes in the magnetic fields. In a previous analysis, Mayfield et al. [1970] have discussed in detail the relation between millimeter enhancement and magnetic fields.




Figure 1. The 3.3-mm solar radio temperature contour map made on 30 June 1971 superposed on a concurrent H α photograph. The contours are labeled in percent enhancement relative to the temperature of the undisturbed regions denoted by the hatched contours.

SUMMARY OF RESULTS

During the 4 years of observations (from 1970 through 1973) a synoptic presentation was used based on the Zürich heliographic maps of the Sun by Waldmeier [1968] and on the magnetic field presentation of Mt. Wilson by Howard. These maps are arranged to present the 360° of solar longitude, showing the millimeter isotherms on a day near central meridian passage. A typical synoptic map is shown in Figure 2 for Carrington rotation 1521. The Carrington longitude is given at the top of the figure, and the date of central meridian passage is located by the filled circle and day number at the bottom. Solar latitude between $\pm 50^\circ$ forms the vertical scale. Maps used in the synoptic presentation have been normalized to the quiet regions indicated by the crosses in the figure; times at which the maps were obtained are indicated with tic marks at the bottom of the figure. For presentation in this synoptic format, only isotherms representing 4% or more enhancement have been used, because they identify enhanced regions and show the close correlation between the millimeter enhancement and white light faculae evident in the Zürich maps.

Also plotted in Figure 2 is an overlay of sunspots and faculae taken from the Zürich map for rotation 1521. This shows a close correlation between the 4% isotherm contour and areas of faculae in sunspot regions. An interpretation of millimeter emission and its relation to white light faculae has been given by Mayfield et al. [1973] and will not be discussed here. Synoptic maps for 1970 through 1973 are given in the appendix.

Synoptic maps for 1970 through 1973 are given in the appendix. They have been plotted to facilitate comparison with the Zürich photoheliographic maps and with the synoptic magnetic field maps published in the IAU's *Quarterly Bulletin of Solar Activity*.




Figure 2. Synoptic 3.3-mm radio temperature contour map for Carrington rotation 1521 superposed on the corresponding heliographic map of the photosphere by Waldmeier [1968]. Carrington Longitude appears at the top of the figure and date of CMP is given at the bottom. The tic marks along the bottom margin of the rectangular grid indicate times at which the maps were taken. Vertical scale is solar latitude. The contours are at levels of 4, 6, 8%, etc. enhancement above the quiet regions indicated by crosses. Peak enhancements are shown by the filled circles. Note the close correlation between the 4% contour levels and facular areas and between peak enhancements and sunspots.

The longevity of active regions can be investigated by inspecting successive synoptic maps and noting the recurrence of active regions. In this respect it is important to point out the sensitivity and precision of the data contained in the synoptic maps, because these parameters influence the visibility of the enhancements. The lowest contour level plotted corresponds to a 4% enhancement, so chosen because individual active regions at this level are usually discernible. A 3% contour, on the other hand, typically runs nearly the length of the map, providing very little information. Additionally, measurements made on different daily maps, either on the same day or on different days, can vary as much as $\pm 0.5\%$. The average, though, probably lies closer to $\pm 0.3\%$; some outstanding examples show variations of no more than $\pm 0.2\%$ over a 5- or 6-day period. To maintain a precision of about 10% (e.g., $4 \pm 0.4\%$), it was safest to display the 4% contour as the lowest level.

Therefore, the longevity of an active region is defined as the time it first exceeds the 4% enhancement level to the time it first drops below that level. Based on this criterion, some regions are found to last less than a full solar rotation; others can persist and be followed over 5 or 6 rotations. As will be shown below, this result is entirely consistent with the strong correlation between the shapes and locations of the 4% contour levels and the outlines of the facular areas--areas that can have lifetimes as short as a few days or as long as a few solar rotations.

As indicated by the clustering of crosses, quiet regions represent another feature of the synoptic maps, the longevity of which can be gauged. Certain examples of these regions can be followed over 3 or 4 rotations even though their nature remains in question.

Daily observations of sunspots and photospheric faculae have been presented in synoptic form by Waldmeier [1968] in *Heliographic Maps of the Photosphere*. An example of these maps (Figure 2) shows each sunspot group at the time of its maximum evolution. In the maps umbrae are depicted by black dots, penumbrae by their outlines, and faculae by dashed lines.

Comparison of the radio synoptic map with the heliographic map in Figure 2 shows how well the 4% enhancement contours correspond to the extensions of the facular regions. Differences in shape and extent can be attributed to a number of causes that include (1) slow, evolutionary changes in the active regions occurring between the maximum phase captured in the photospheric maps and the stage nearest central meridian passage reflected in the millimeter maps; (2) flare-induced rapid changes in the active regions, the residual effects of which can influence the radio contours (although, such effects have been allowed for and corrected whenever possible); (3) selection of an invalid normalization point for a particular daily radio map such as a point containing a dark absorption filament, which would cause an apparent "growth" or "contraction" of the enhancement levels; and (4) lack of a physical connection between the millimeter emission and the white light faculae and sunspots.

ACKNOWLEDGMENTS



This research was supported in part by company funds of The Aerospace Corporation.



REFERENCES



- | | | |
|---|------|--|
| JACOBS, E. and
H.E. KING | 1965 | "2.8 minute beamwidth, millimeter-wave antenna
measurement and evaluation," <i>IEEE International
Convention Record</i> , Part 5, 92-100. |
| KING, H.E.,
E. JACOBS and
J.M. STACEY | 1966 | "A 2.8 arc-min beamwidth antenna: lunar eclipse
observations at 3.2 mm," <i>IEEE Trans. on Antennas
and Propagation</i> , AP-14, 82-91. |
| MAYFIELD, E.B.,
J. HIGMAN and
C. SAMSON | 1970 | "Variations in solar emission at 3.3 mm wavelength
and their relation to flares," <i>Solar Phys.</i> , 13,
372-388. |
| MAYFIELD, E.B.,
K.P. WHITE III and
F.I. SHIMABUKURO | 1973 | "Millimeter radio evidence for containment
mechanisms in solar flares," in <i>Coronal Disturbances</i> ,
IAU Symposium 57, Gorden Newkirk, Jr., Ed. (D. Riedel
Publishing Company, Dordrecht), 447-460. |
| MAYFIELD, E.B.,
K.P. WHITE III and
F.I. SHIMABUKURO | 1974 | "Synoptic radio maps of the sun at 3.3 mm for the
years 1967-1969," <i>Report UAG-32</i> , (Department of
Commerce, NOAA/NGSDC, Boulder, CO). |
| SHIMABUKURO, F.I. | 1968 | "A comparison of 3.3-mm bursts and H α emission during
flares," <i>Solar Phys.</i> , 5, 498-506. |
| WALDMEIER, M. | 1968 | "Heliographic maps of the photosphere for the year
1967," <i>Publikationen der Eidgenössischen Sternwarte
Zürich</i> , 13, No. 2. |



APPENDIX



Synoptic Radio Maps of the Sun at 3.3 mm, 1970-73



























































































UAG Series of Reports

Prepared by World Data Center A for Solar-Terrestrial Physics, NOAA, Boulder, Colorado, U.S.A.

These reports are for sale through the National Climatic Center, Federal Building, Asheville, NC 28801, Attn: Publications. Subscription price: \$25.20 a year; \$12.00 additional for foreign mailing; single copy price varies. These reports are issued on an irregular basis with 6 to 12 reports being issued each year. Therefore, in some years the single copy rate will be less than the subscription price, and in some years the single copy rate will be more than the subscription price. Make check or money order payable to: Department of Commerce, NOAA.

Some issues are now out of print and are available only on microfiche as indicated. Requests for microfiche should be sent to World Data Center A for Solar-Terrestrial Physics, NOAA, Boulder, CO 80302, with check or money order made payable to Department of Commerce, NOAA.

- UAG-1 "IQSY Night Airglow Data", price \$1.75.
UAG-2 "A Reevaluation of Solar Flares, 1964-1966", price 30 cents.
UAG-3 "Observations of Jupiter's Sporadic Radio Emission in the Range 7.6-41 MHz, 6 July 1966 through 8 September 1968", microfiche only, price 45 cents.
UAG-4 "Abbreviated Calendar Record 1966-1967", price \$1.25.
UAG-5 "Data on Solar Event of May 23, 1967 and its Geophysical Effects", price 65 cents.
UAG-6 "International Geophysical Calendars 1957-1969", price 30 cents.
UAG-7 "Observations of the Solar Electron Corona: February 1964-January 1968", price 15 cents.
UAG-8 "Data on Solar-Geophysical Activity October 24-November 6, 1968", price (includes Parts 1 & 2) \$1.75.
UAG-9 "Data on Cosmic Ray Event of November 18, 1968 and Associated Phenomena", price 55 cents.
UAG-10 "Atlas of Ionograms", price \$1.50.
UAG-11 "Catalogue of Data on Solar-Terrestrial Physics" (now obsolete).
UAG-12 "Solar-Geophysical Activity Associated with the Major Geomagnetic Storm of March 8, 1970", price (includes Parts 1-3) \$3.00.
UAG-13 "Data on the Solar Proton Event of November 2, 1969 through the Geomagnetic Storm of November 8-10, 1969, price 50 cents.
UAG-14 "An Experimental, Comprehensive Flare Index and Its Derivation for 'Major' Flares, 1955-1969", price 30 cents.
UAG-15 "Catalogue of Data on Solar-Terrestrial Physics" (now obsolete).
UAG-16 "Temporal Development of the Geographical Distribution of Auroral Absorption for 30 Substorm Events in each of IQSY (1964-65) and IASY (1969)", price 70 cents.
UAG-17 "Ionospheric Drift Velocity Measurements at Jicamarca, Peru (July 1967-March 1970)", microfiche only, price 45 cents.
UAG-18 "A Study of Polar Cap and Auroral Zone Magnetic Variations", price 20 cents.
UAG-19 "Reevaluation of Solar Flares 1967", price 15 cents.
UAG-20 "Catalogue of Data on Solar-Terrestrial Physics" (now obsolete).
UAG-21 "Preliminary Compilation of Data for Retrospective World Interval July 26 - August 14, 1972", price 70 cents.
UAG-22 "Auroral Electrojet Magnetic Activity Indices (AE) for 1970", price 75 cents.
UAG-23 "U.R.S.I. Handbook of Ionogram Interpretation and Reduction", price \$1.75.
UAG-24 "Data on Solar-Geophysical Activity Associated with the Major Ground Level Cosmic Ray Events of 24 January and 1 September 1971", price (includes Parts 1 and 2) \$2.00.
UAG-25 "Observations of Jupiter's Sporadic Radio Emission in the Range 7.6-41 MHz, 9 September 1968 through 9 December 1971", price 35 cents.
UAG-26 "Data Compilation for the Magnetospherically Quiet Periods February 19-23 and November 29 - December 3, 1970", price 70 cents.
UAG-27 "High Speed Streams in the Solar Wind", price 15 cents.
UAG-28 "Collected Data Reports on August 1972 Solar-Terrestrial Events", price (includes Parts 1-3) \$4.50.
UAG-29 "Auroral Electrojet Magnetic Activity Indices AE (11) for 1968", price 75 cents.
UAG-30 "Catalogue of Data on Solar-Terrestrial Physics", price \$1.75.
UAG-31 "Auroral Electrojet Magnetic Activity Indices AE (11) for 1969", price 75 cents.
UAG-32 "Synoptic Radio Maps of the Sun at 3.3 mm for the Years 1967-1969", price 35 cents.
UAG-33 "Auroral Electrojet Magnetic Activity Indices AE (10) for 1967", price 75 cents.
UAG-34 "Absorption Data for the IGY/IGC and IQSY", price \$2.00.
UAG-35 "Catalogue of Digital Geomagnetic Variation Data at World Data Center A for Solar-Terrestrial Physics", price 20 cents.
UAG-36 "An Atlas of Extreme Ultraviolet Flashes of Solar Flares Observed Via Sudden Frequency Deviations During the ATM-SKYLAB Missions", price 55 cents.
UAG-37 "Auroral Electrojet Magnetic Activity Indices AE (10) for 1966", price 75 cents.
UAG-38 "Master Station List for Solar-Terrestrial Physics Data at WDC-A for Solar-Terrestrial Physics", price \$1.60.
UAG-39 "Auroral Electrojet Magnetic Activity Indices AE (11) for 1971", by Joe Haskell Allen, Carl C. Abston and Leslie D. Morris, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, February 1975, 144 pages, price \$2.05.
UAG-40 "H-Alpha Synoptic Charts of Solar Activity For the Period of Skylab Observations, May, 1973-March, 1974", by Patrick S. McIntosh, NOAA Environmental Research Laboratories, February 1975, 32 pages, price 56 cents.
UAG-41 "H-Alpha Synoptic Charts of Solar Activity During the First Year of Solar Cycle 20, October, 1964 - August, 1965", by Patrick S. McIntosh, NOAA Environmental Research Laboratories, and Jerome T. Nolte, American Science and Engineering, Cambridge, Massachusetts, March 1975, 25 pages, price 48 cents.
UAG-42 "Observations of Jupiter's Sporadic Radio Emission in the Range 7.6-80 MHz 10 December 1971 through 21 March 1975", by James W. Warwick, George A. Dulk, and Anthony C. Riddle, Department of Astro-Geophysics, University of Colorado, Boulder, Colorado 80302, April 1975, 49 pages, price \$1.15.
UAG-43 "Catalog of Observation Times of Ground-Based Skylab-Coordinated Solar Observing Programs", compiled by Helen E. Coffey, World Data Center A for Solar-Terrestrial Physics, May 1975, 159 pages, price \$3.00.
UAG-44 "Synoptic Maps of Solar 9.1 cm Microwave Emission from June 1962 to August 1973", by Werner Graf and Ronald N. Bracewell, Radio Astronomy Institute, Stanford University, Stanford, California 94305, May 1975, 183 pages, price \$2.55.
UAG-45 "Auroral Electrojet Magnetic Activity Indices AE (11) for 1972", by Joe Haskell Allen, Carl C. Abston and Leslie D. Morris, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, May 1975, 144 pages, price \$2.10.
UAG-46 "Interplanetary Magnetic Field Data 1963-1974", by Joseph H. King, National Space Science Data Center, NASA Goddard Space Flight Center, Greenbelt, Maryland 20771, June 1975, 382 pages, price \$2.95.
UAG-47 "Auroral Electrojet Magnetic Activity Indices AE (11) for 1973", by Joe Haskell Allen, Carl C. Abston and Leslie D. Morris, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, June 1975, 144 pages, price \$2.10.

- UAG-48A "Synoptic Observations of the Solar Corona during Carrington Rotations 1580-1596 (11 October 1971 - 15 January 1973)", [Reissue with quality images] by R. A. Howard, M. J. Koomen, D. J. Michels, R. Tousey, C. R. Detwiler, D. E. Roberts, R. T. Seal and J. D. Whitney, E. O. Hulbert Center for Space Research, NRL, Washington, D. C. 20375 and R. T. and S. F. Hansen, C. J. Garcia and E. Yasukawa, High Altitude Observatory, NCAR, Boulder, Colorado 80303, February 1976, 200 pages, price \$4.27.
- UAG-49 "Catalog of Standard Geomagnetic Variation Data", prepared by Environmental Data Service, NOAA, Boulder, Colorado, August 1975, 125 pages, price \$1.85.
- UAG-50 "High-Latitude Supplement to the URSI Handbook on Ionogram Interpretation and Reduction", by W. R. Piggott, British Antarctic Survey, c/o SRC, Appleton Laboratory, Ditton Park, Slough, England, October 1975, 292 pages, price \$4.00.
- UAG-51 "Synoptic Maps of Solar Coronal Hole Boundaries Derived from He II 304Å Spectroheliograms from the Manned Skylab Missions", by J. D. Bohlin and D. M. Rubenstein, E. O. Hulbert Center for Space Research, Naval Research Observatory, Washington, D. C. 20375 U.S.A., November 1975, 30 pages, price 54 cents.
- UAG-52 "Experimental Comprehensive Solar Flare Indices for Certain Flares, 1970-1974", compiled by Helen W. Dodson and E. Ruth Hedeman, McMath-Hulbert Observatory, The University of Michigan, 895 Lake Angelus Road North, Pontiac, Michigan 48055 U.S.A., November 1975, 27 pages, price 60 cents.
- UAG-53 "Description and Catalog of Ionospheric F-Region Data, Jicamarca Radar Observatory (November 1966 - April 1969)", by W. L. Clark and T. E. Van Zandt, Aeronomy Laboratory, NOAA, Boulder, Colorado 80302 and J. P. McClure, University of Texas at Dallas, Dallas, Texas 75230, April 1976, 10 pages, price 33 cents.
- UAG-54 "Catalog of Ionosphere Vertical Soundings Data", prepared by Environmental Data Service, NOAA, Boulder, Colorado 80302, April 1976, 130 pages, price \$2.10.
- UAG-55 "Equivalent Ionospheric Current Representations by a New Method, Illustrated for 8-9 November 1969 Magnetic Disturbances", by Y. Kamide, Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, Colorado 80302 and Geophysical Institute, University of Alaska, Fairbanks, Alaska 99701, H. W. Kroehl, Data Studies Division, NOAA/EDS/NGSDC, Boulder, Colorado 80302, M. Kanamitsu, Advanced Study Program, National Center for Atmospheric Research, Boulder, Colorado 80303, J. H. Allen, Data Studies Division, NOAA/EDS/NGSDC, Boulder, Colorado 80302, and S.-I. Akasofu, Geophysical Institute, University of Alaska, Fairbanks, Alaska 99701, April 1976, 91 pages, price \$1.60.
- UAG-56 "Iso-intensity Contours of Ground Magnetic H Perturbations for the December 16-18, 1971 Geomagnetic Storm", by Y. Kamide, Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, Colorado 80302 and Geophysical Institute, University of Alaska, Fairbanks, Alaska 99701 (currently Guest worker at Data Studies Division, NOAA/EDS/NGSDC, Boulder, Colorado 80302), April 1976, 37 pages, price \$1.39.
- UAG-57 "Manual on Ionospheric Absorption Measurements", edited by K. Rawer, Institut für Physikalische Weltraumforschung, Freiburg, G.F.R., June 1976, 202 pages, price \$4.27.
- UAG-58 "ATS6 Radio Beacon Electron Content Measurements at Boulder, July 1974 - May 1975", by R. B. Fritz, Space Environment Laboratory (currently with Wave Propagation Laboratory), NOAA, Boulder, Colorado 80302 USA, September 1976, 61 pages, price \$1.04.
- UAG-59 "Auroral Electrojet Magnetic Activity Indices AE(11) for 1974", by Joe Haskell Allen, Carl C. Abston and Leslie D. Morris, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, December 1976, 144 pages, price \$2.16.
- UAG-60 "Geomagnetic Data for January 1976 (AE(7) Indices and Stacked Magnetograms)" by J. H. Allen, C. A. Abston and L. R. Morris, NGSDC/EDS/NOAA, July 1977, 57 pages, price \$1.07.
- UAG-61 "Collected Data Reports for STIP Interval II 20 March - 5 May 1976", edited by Helen E. Coffey and John A. McKinnon, National Geophysical and Solar-Terrestrial Data Center, Environmental Data Service, August 1977, 313 pages, price \$2.95.
- UAG-62 "Geomagnetic Data for February 1976 (AE(7) Indices and Stacked Magnetograms)" by J. H. Allen, C. A. Abston and L. R. Morris, NGSDC/EDS/NOAA, September 1977, 55 pages, price \$1.11.
- UAG-63 "Geomagnetic Data for March 1976 (AE(7) Indices and Stacked Magnetograms)" by J. H. Allen, C. A. Abston and L. R. Morris, NGSDC/EDS/NOAA, September 1977, 57 pages, price \$1.11
- UAG-64 "Geomagnetic Data for April 1976 (AE(8) Indices and Stacked Magnetograms)" by J. H. Allen, C. A. Abston and L. R. Morris, NGSDC/EDS/NOAA, February 1978.